

A Fubini Counterexample

Fubini's Theorem states

Theorem (Fubini) *If $f(x, y)$ is continuous in a region R described by both*

$$x_1 \leq x \leq x_2 \quad y_1(x) \leq y \leq y_2(x)$$

and

$$y_1 \leq y \leq y_2 \quad x_1(y) \leq x \leq x_2(y)$$

with $y_1(x)$, $y_2(x)$, $x_1(y)$ and $x_2(y)$ continuous, then

$$\int_{x_1}^{x_2} dx \left[\int_{y_1(x)}^{y_2(x)} dy f(x, y) \right] \quad \text{and} \quad \int_{y_1}^{y_2} dy \left[\int_{x_1(y)}^{x_2(y)} dx f(x, y) \right]$$

both exist and are equal.


In these notes, we relax exactly one of the hypotheses of Fubini's Theorem, namely the continuity of f , and construct an example in which both of the integrals in Fubini's Theorem exist, but are **not equal**. In fact, we choose $R = \{ (x, y) \mid 0 \leq x \leq 1, 0 \leq y \leq 1 \}$ and we use a function $f(x, y)$ that is continuous on R , except at exactly one point – the origin.

First, let $\delta_1, \delta_2, \delta_3, \dots$ be any sequence of real numbers obeying

$$1 = \delta_1 > \delta_2 > \delta_3, \dots, \delta_n \rightarrow 0$$

For example $\delta_n = \frac{1}{n}$ or $\delta_n = \frac{1}{2^{n-1}}$ are both acceptable. For each positive integer n , let $I_n = (\delta_{n+1}, \delta_n] = \{ t \mid \delta_{n+1} < t \leq \delta_n \}$ and let $g_n(t)$ be any continuous function obeying $g_n(\delta_{n+1}) = g_n(\delta_n) = 0$ and $\int_{I_n} g(t) dt = 1$. There are many such functions. For example


$$g_n(t) = \frac{2}{\delta_n - \delta_{n+1}} \begin{cases} \delta_n - t & \text{if } \frac{1}{2}(\delta_{n+1} + \delta_n) \leq t \leq \delta_n \\ t - \delta_{n+1} & \text{if } \delta_{n+1} \leq t \leq \frac{1}{2}(\delta_{n+1} + \delta_n) \\ 0 & \text{otherwise} \end{cases}$$



Now define

$$f(x, y) = \begin{cases} 0 & \text{if } x = 0 \\ 0 & \text{if } y = 0 \\ g_m(x)g_n(y) & \text{if } x \in I_m, y \in I_n \text{ with } m = n \\ -g_m(x)g_n(y) & \text{if } x \in I_m, y \in I_n \text{ with } m = n + 1 \\ 0 & \text{otherwise} \end{cases}$$

You should think of $(0, 1] \times (0, 1]$ as a union of a bunch of small rectangles $I_m \times I_n$, as in the figure below. On most of these rectangles, $f(x, y)$ is just zero. The exceptions are the darkly shaded rectangles $I_n \times I_n$ on the “diagonal” of the figure and the lightly shaded rectangles $I_{n+1} \times I_n$ just to the left of the “diagonal”. On each darkly shaded rectangle, the graph of $f(x, y)$ is the graph of $g_n(x)g_n(y)$ which looks like a pyramid. On each lightly shaded rectangle, the graph of $f(x, y)$ is the graph of $-g_{n+1}(x)g_n(y)$ which looks like a pyramidal hole in the ground.



Now fix any $0 \leq y \leq 1$ and let's compute $\int_0^1 f(x, y) dx$. That is, we are integrating f along a line that is parallel to the x -axis. If $y = 0$, then $f(x, y) = 0$ for all x , so $\int_0^1 f(x, y) dx = 0$. If $0 < y \leq 1$, then there is exactly one positive integer n with $y \in I_n$ and $f(x, y)$ is zero, except for x in I_n or I_{n+1} . So for $y \in I_n$

$$\begin{aligned} \int_0^1 f(x, y) dx &= \sum_{m=n, n+1} \int_{I_m} f(x, y) dx = \int_{I_n} g_n(x)g_n(y) dx - \int_{I_{n+1}} g_{n+1}(x)g_n(y) dx \\ &= g_n(y) \int_{I_n} g_n(x) dx - g_n(y) \int_{I_{n+1}} g_{n+1}(x) dx \\ &= g_n(y) - g_n(y) = 0 \end{aligned}$$

Here we have twice used that $\int_{I_m} g(t) dt = 1$ for all m . Thus $\int_0^1 f(x, y) dx = 0$ for all y and hence $\int_0^1 dy \left[\int_0^1 dx f(x, y) \right] = 0$.

Finally, fix any $0 \leq x \leq 1$ and let's compute $\int_0^1 f(x, y) dy$. That is, we are integrating f along a line that is parallel to the y -axis. If $x = 0$, then $f(x, y) = 0$ for all y , so $\int_0^1 f(x, y) dy = 0$. If $0 < x \leq 1$, then there is exactly one positive integer m with $x \in I_m$. If $m \geq 2$, then $f(x, y)$ is zero, except for y in I_m and I_{m-1} . But, if $m = 1$, then $f(x, y)$ is zero, except for y in I_1 . (Take another look at the figure on the previous page.) So for $x \in I_m$, with $m \geq 2$,

$$\begin{aligned} \int_0^1 f(x, y) dy &= \sum_{n=m, m-1} \int_{I_n} f(x, y) dy = \int_{I_m} g_m(x)g_m(y) dy - \int_{I_{m-1}} g_m(x)g_{m-1}(y) dy \\ &= g_m(x) \int_{I_m} g_m(y) dy - g_m(x) \int_{I_{m-1}} g_{m-1}(y) dy \\ &= g_m(x) - g_m(x) = 0 \end{aligned}$$

But $x \in I_1$,

$$\int_0^1 f(x, y) dy = \int_{I_1} f(x, y) dy = \int_{I_1} g_1(x)g_1(y) dy = g_1(x) \int_{I_1} g_1(y) dy = g_1(x)$$

Thus

$$\int_0^1 f(x, y) dy = \begin{cases} 0 & \text{if } x \leq \delta_2 \\ g_1(x) & \text{if } x \in I_1 \end{cases}$$

and hence

$$\int_0^1 dx \left[\int_0^1 dy f(x, y) \right] = \int_{I_1} g_1(x) dx = 1$$

The conclusion is that for the $f(x, y)$ above, which is defined for all $0 \leq x \leq 1$ and $0 \leq y \leq 1$ and is continuous except at $(0, 0)$,

$$\int_0^1 dy \left[\int_0^1 dx f(x, y) \right] = 0 \quad \int_0^1 dx \left[\int_0^1 dy f(x, y) \right] = 1$$